


Name

L.I.U.

Chem. 122, Sect 007,

Quiz 1, 50 pts, Spring, 2012

1. For the following molecules identify (a) the number of proton signals, their splitting patterns and their relative intensities (b) the number of carbon signals. (15 pts)



2. Identify the following molecule of formula C₁₀H₁₁ClO₂; IR: 1735 cm⁻¹; ¹H: δ 1.2, triplet, 3H; 2.8, singlet, 2H; 3.9, quartet, 2H; 7.2, doublet, 2H; 7.4, doublet, 2H. (15 pts)

3. Answer the following questions on the preparation of acetyl salicylic acid from salicylic acid, acetic anhydride and phosphoric acid. (a) Is salicylic acid soluble in acetic anhydride? Explain briefly. (b) Why is it important to keep the spatula in the test tube at all times during recrystallization? (c) One student used a large test tube during the recrystallization instead of a small test tube and ended up adding about 20 mL of the 50:50 ethyl acetate/hexane solvent. All of her solid dissolved in the hot solution but when she cooled it down to room temperature and then in an ice bath she did not get any crystals formed. Was her experiment ruined? What could she do to recover her product? (7 pts)

4. In the preparation of triphenyl carbinol from bromobenzene, magnesium and methyl benzoate (C₆H₅CO₂CH₃) (a) show the overall reaction, including all intermediates. (b) What is the purpose of the drying tube? (c) How can you tell that you have formed the organomagnesium (Grignard) reagent? (d) What is the purpose of adding the aqueous HCl at the end of the reaction? Show the reaction that occurs. (e) Do you need to have the drying tube attached to your reaction flask at this point? Explain briefly. (13 pts)